

Treatment and rehabilitation of patellar tendinitis

Jinyang Wang

Nanjing University of Chinese Medicine, Nanjing, China

wangjinyang@ldy.edu.rs

Abstract. Patellar tendinitis, commonly known as "jumper's knee", is a prevalent overuse injury that primarily affects athletes involved in sports requiring frequent jumping and abrupt directional changes. This review provides a comprehensive analysis of the pathophysiology, causes, and injury mechanisms associated with patellar tendinitis, focusing on the role of repetitive mechanical overload and circulatory disturbances at the tendon's insertional region. The high tensile stress experienced during jumping and landing activities significantly contributes to tendon microtears and degeneration, potentially leading to chronic tendinopathy if left untreated. The review also classifies patellar tendinitis based on severity, ranging from mild pain to complete tendon rupture. Both surgical and conservative treatment options are discussed, with surgical intervention recommended for severe cases and conservative approaches such as pharmacotherapy, eccentric strength training, patellar taping, and shockwave therapy shown to be effective in reducing pain and promoting tendon healing. This review aims to provide valuable insights for clinicians and sports professionals, enhancing understanding of injury mechanisms and treatment strategies to improve patient outcomes and athletic performance.

Keywords: Patellar tendinitis, Mechanical overload, Circulatory disturbances, Conservative treatment, Eccentric strength training.

1. Introduction

Patellar tendinitis, often referred to as patellar ligament inflammation or "jumper's knee," is a prevalent overuse injury predominantly affecting athletes involved in sports characterized by frequent jumping, abrupt directional changes, and rapid deceleration [1]. This condition is marked by chronic inflammation of the patellar tendon, typically resulting from repetitive mechanical stress and inadequate recovery. Patellar tendinitis falls under the category of overuse syndromes, where persistent mechanical stress during activities such as jumping and sudden stops exceeds the tendon's ability to repair itself, leading to progressive damage [2].

The pathophysiology of patellar tendinitis involves microtears in the collagen fibers of the tendon due to excessive tensile stress. These microtears accumulate over time, resulting in degenerative changes such as disorganized collagen fibers, increased cellularity, and neovascularization. The chronic inflammatory response and subsequent tendon degeneration contribute to the pain and functional limitations experienced by affected individuals. Without appropriate management, patellar tendinitis can progress to severe tendon degeneration, which can significantly impair athletic performance and cause long-term damage [3].

Athletes participating in sports with repetitive jumping and landing, such as volleyball, basketball, and soccer, are particularly at risk. Epidemiological studies suggest that approximately 40% of volleyball players suffer from patellar tendinitis, underscoring the high-risk nature of these high-impact sports. The condition is also prevalent among high-level basketball and soccer players, highlighting the need for early recognition and intervention to address tendon stress and inflammation effectively [4]. Patellar tendinitis not only impacts athletic performance but also poses significant challenges to an athlete's overall health and career longevity. The condition typically presents as pain localized to the anterior aspect of the knee, especially around the lower pole of the patella. Pain usually exacerbates with activity, particularly during jumping or squatting, and may persist at rest in advanced cases. Early diagnosis and intervention are crucial to prevent further tendon damage and to facilitate a successful return to competitive activities [5].

In summary, patellar tendinitis is a significant overuse injury common among athletes involved in sports with frequent jumping and rapid deceleration. Understanding the risk factors, pathophysiology, and clinical implications of this condition is essential for developing effective treatment and rehabilitation strategies, which are critical for maintaining athletic performance and preventing long-term complications.

2. Injury mechanisms of patellar tendinitis

Athletic activities that involve frequent jumping, rapid deceleration, and sudden direction changes—such as basketball, volleyball, and soccer—are particularly prone to causing patellar tendinitis. This condition, primarily affects athletes due to the significant mechanical stress placed on the patellar tendon during these movements. The exact etiology of patellar tendinitis remains complex and not fully understood; however, it can be broadly categorized into two main aspects: local mechanical overload on the patellar tendon and circulatory disturbances in the tendon's insertional region. Understanding these injury mechanisms is essential for developing effective prevention and treatment strategies.

2.1. Local mechanical overload on the patellar tendon

From a biomechanical perspective, the patellar tendon is essential for transmitting the force generated by the quadriceps muscle during various physical activities. When the quadriceps contract, the resulting force is transmitted through the patellar tendon to the tibia, facilitating movements such as jumping, running, and landing. This process subjects the tendon to significant tensile stress. Under normal conditions, the patellar tendon can withstand substantial forces. However, when the tensile load exceeds the tendon's physiological capacity or is applied repeatedly without sufficient recovery, pathological changes can occur.

Mechanical overload on the patellar tendon can cause microtears in the collagen fibers, initiating a cycle of injury and repair that may lead to tendinopathy. Repeated exposure to excessive tensile stress can overwhelm the tendon's ability to repair itself, resulting in the gradual accumulation of microtears, inflammation, and ultimately, tendon degeneration. This issue is particularly pronounced in athletes involved in sports with frequent jumping and sudden changes in direction, such as volleyball, basketball, and soccer. In these sports, the patellar tendon may endure forces up to seven times the athlete's body weight during sudden stops and jumps. Similarly, during the landing phase, the tendon can experience tensile forces of up to 5.5 times the body weight [5]. Such substantial loads place significant strain on the patellar tendon, making it highly susceptible to injury.

In younger populations, particularly adolescents, the patellar tendon is still developing and may be more prone to injury due to the intense mechanical stress associated with sports activities. The repetitive pulling forces exerted on the tendon can lead to the development of tibial tubercle osteochondrosis, commonly known as Osgood-Schlatter disease. This condition involves inflammation and pain at the growth plate of the tibial tubercle and is prevalent among young athletes participating in high-impact sports. The mechanism behind this condition involves repeated stress on the growth plate by the patellar tendon, potentially causing microtears, inflammation, and, in severe cases, fragmentation of the tibial tubercle [6].

Overall, local mechanical overload is a critical factor in the development of patellar tendinitis. Excessive tensile stress during physical activity can lead to microtears, inflammation, and degeneration of the patellar tendon, ultimately resulting in tendinopathy. This problem is especially evident in athletes who engage in sports requiring frequent jumping, rapid deceleration, and abrupt changes in direction, rendering them highly susceptible to patellar tendinitis.

2.2. Local circulatory disturbances in the tendon's insertional region

In addition to mechanical overload, circulatory disturbances in the insertional region of the patellar tendon have been identified as significant contributors to the development of patellar tendinitis. Anatomical studies and angiographic examinations show that the blood supply to the patellar tendon, particularly at its insertional region, is relatively poor. This area, located near the tendon's attachment to the patella, possesses a limited vascular network, making it more vulnerable to ischemia and subsequent tissue damage [7].

During activities involving repetitive jumping, running, and landing, the patellar tendon experiences continuous tensile stress, which can compromise blood flow to the insertional region. The sustained mechanical loading increases intratendinous pressure, particularly within the fibrocartilaginous zones, which are already poorly vascularized. This increased pressure impairs the diffusion of essential nutrients and oxygen to tendon cells, leading to localized ischemia and subsequent tissue hypoxia [8]. Consequently, the tendon's ability to repair itself is compromised, leading to the accumulation of metabolic waste products and the initiation of an inflammatory response.

Chronic overuse of the patellar tendon, combined with impaired blood circulation, can cause microtrauma within the tendon's insertional region. Over time, these microtraumas result in degenerative changes, including collagen disorganization, increased cellularity, and neovascularization. While neovascularization is initially a reparative response, it can exacerbate the condition by introducing fragile blood vessels prone to rupture. The resultant microhemorrhages further compromise the local blood supply, creating a vicious cycle of ischemia, inflammation, and tendon degeneration [9].

Chronic circulatory disturbances also contribute to the development of tendinopathy by promoting fibrotic tissue formation. Repeated episodes of ischemia and inflammation stimulate the production of fibrotic tissue, which replaces normal tendon tissue. This fibrotic tissue is less elastic and more prone to injury, further compromising the tendon's function and increasing the risk of rupture. In severe cases, prolonged ischemia and inflammation can lead to tissue necrosis and avascular necrosis, characterized by tendon tissue death due to a lack of blood supply [10].

Circulatory disturbances in the patellar tendon's insertional region are further exacerbated by the increased metabolic demands placed on the tendon during intense physical activity. As tendon cells become more metabolically active, their demand for oxygen and nutrients rises. However, the compromised blood supply cannot meet these demands, leading to additional tissue damage and inflammation [11]. This imbalance between the tendon's metabolic needs and the available blood supply perpetuates chronic ischemia, further damaging and inflaming the tissue.

In conclusion, the development of patellar tendinitis is a multifactorial process involving both local mechanical overload and circulatory disturbances in the tendon's insertional region. Excessive tensile stress during physical activity leads to microtears, inflammation, and degeneration of the patellar tendon. At the same time, impaired blood circulation exacerbates these pathological changes. Understanding these mechanisms is essential for developing effective prevention and treatment strategies, particularly for athletes at high risk for this condition.

3. Classification of patellar tendinitis injuries

The grading of patellar tendinopathy is typically based on the severity of symptoms, the extent of functional limitations, and imaging findings. In Grade I, patients experience mild pain during or after activity without impairment of normal function; in Grade II, pain intensifies during or after activity with mild functional restrictions; in Grade III, severe and persistent pain significantly limits function, and imaging shows widespread tendon pathology; Grade IV involves complete tendon rupture, leading to

loss of knee joint function and necessitating surgical repair. Based on these gradings, clinicians can develop appropriate treatment and rehabilitation plans [12].

4. Rehabilitation and treatment of patellar tendinopathy

The rehabilitation and treatment of patellar tendinopathy involve a combination of surgical intervention and conservative management. The choice of treatment depends on the severity of the condition and the effectiveness of conservative approaches. Given the chronic nature of patellar tendinopathy, a comprehensive and often extended rehabilitation process is necessary.

4.1. Surgical treatment

Surgical intervention is typically considered when patellar tendinopathy reaches Grade IIIb or IV, or when conservative treatments fail to produce significant improvement. The primary goal of surgery is to remove damaged tissue and restore the patellar tendon's normal structure and function. Common surgical techniques include patellar tendon release, tendon repair, and excision of diseased tissue [8]. Post-surgery, patients require extensive rehabilitation to promote tissue healing and functional recovery.

4.2. Conservative treatment

Conservative treatment is essential for managing patellar tendinopathy, especially given its prolonged course. It focuses on reducing knee joint load, avoiding excessive physical activity, and allowing adequate time for tendon repair. Conservative methods include pharmacotherapy, physical factor therapy, patellar taping, and strength training [13]. These methods not only help alleviate symptoms but also improve the healing environment of the patellar tendon, preventing further deterioration.

4.2.1. Pharmacotherapy. Pharmacotherapy is primarily used for pain relief in patellar tendinopathy. Non-steroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen and naproxen, are commonly employed to alleviate pain. Corticosteroid injections or medications can also provide short-term relief. However, while these medications are effective in managing acute symptoms, they do not address the underlying tendon pathology and may have side effects with prolonged use. Thus, pharmacotherapy is usually a short-term solution for symptom control.

4.2.2. Physical factor therapy. Physical factor therapy is a key component of conservative treatment and has been increasingly utilized for patellar tendinopathy. Techniques such as cryotherapy, ultrasound therapy, electrotherapy, traction therapy, and shockwave therapy are effective in reducing pain and inflammation. Cryotherapy lowers local tissue temperature to alleviate pain; ultrasound therapy uses high-frequency vibrations to promote tissue repair; electrotherapy and traction therapy help relieve muscle tension and improve joint mobility; and shockwave therapy stimulates tissue regeneration, aiding recovery. These therapies not only alleviate symptoms but also enhance local blood circulation and support tissue healing.

4.2.3. Patellar taping techniques. Patellar taping is a non-invasive method that provides external support to enhance knee joint function. It improves patellar tendon stability, reduces pain during physical activity, and promotes tissue repair by increasing local blood flow. Studies, such as those by Aminaka, demonstrated the effectiveness of patellar taping in alleviating symptoms of knee joint injuries, including patellar pain syndrome [14]. Using white tape or kinesiology tape to support the patellar tendon can improve local circulation and reduce symptoms.

4.2.4. Strength training. Strength training is crucial for the rehabilitation of patellar tendinopathy, especially in strengthening the muscles surrounding the patellar tendon. Quadriceps atrophy is a significant factor in patellar tendinopathy and other knee joint injuries. Research by Li Haizhen et al. indicated that regular eccentric strength training can significantly reduce pain and improve functional capacity in patients with patellar tendinopathy [14]. Eccentric training increases the tensile capacity of

the patellar tendon, helping it adapt to physical activity and reducing the risk of re-injury. Incorporating other exercises, such as core strengthening and flexibility training, can further enhance athletic performance and support recovery [15].

In summary, a well-designed rehabilitation plan that integrates both surgical and conservative treatments is essential for effectively managing patellar tendinopathy. Such a plan not only alleviates pain and restores knee joint function but also prevents recurrence and supports long-term recovery.

5. Conclusion

This review provides a comprehensive analysis of the mechanisms, causes, and treatment strategies for patellar tendinitis, with a particular focus on athletes engaged in high-impact sports. The research emphasizes the critical roles of mechanical overload and circulatory disturbances within the tendon as primary contributors to the development of this condition. The biomechanical analysis sheds light on how repetitive stress and inadequate recovery contribute to tendon degeneration and inflammation. Additionally, the classification and treatment approaches, encompassing both surgical and conservative methods, provide a structured framework for managing patellar tendinitis according to the severity of the condition.

Looking ahead, the future of patellar tendinitis treatment will likely focus on enhancing early diagnosis, refining rehabilitation protocols, and incorporating innovative therapeutic approaches. Advances in imaging technologies, such as high-resolution ultrasound and MRI, are expected to improve the early detection of tendon degeneration before severe symptoms emerge, enabling more proactive interventions. Furthermore, personalized rehabilitation programs that integrate cutting-edge therapies, including platelet-rich plasma (PRP) injections, stem cell therapy, and advanced shockwave treatments, hold promise for accelerating healing and reducing the recurrence of tendon injuries.

Future research should prioritize a deeper understanding of the cellular and molecular mechanisms underlying tendinitis, particularly in relation to tissue regeneration and repair. The development of targeted pharmacological interventions that address the root causes of tendon degeneration, rather than merely alleviating symptoms, could significantly improve long-term outcomes. By combining these advancements with biomechanical training and injury prevention strategies, the prevalence of patellar tendinitis among athletes could be substantially reduced, leading to quicker and more effective recoveries.

References

- [1] Sheri A. (2005). Etiology of Patellar Tendinopathy in Athletes. *Journal of Sport Rehabilitation*, 3, 259-272.
- [2] Rath, E., Schwarzkopf, R., & Richmond, J. C. (2010). Clinical signs and anatomical correlation of patellar tendinitis. *Indian journal of orthopaedics*, 44(4), 435–437. <https://doi.org/10.4103/0019-5413.69317>
- [3] Bazluki J. (1996). Surgical intervention and rehabilitation of chronic patellar tendinitis. *Journal of athletic training*, 31(1), 65–67.
- [4] Richards, D. P., Ajemian, S. V., Wiley, J. P., & Zernicke, R. F. (1996). Knee joint dynamics predict patellar tendinitis in elite volleyball players. *The American journal of sports medicine*, 24(5), 676–683. <https://doi.org/10.1177/036354659602400520>
- [5] Kotlyar, N., Tsapov, E., Korobeynikov, E., & Deryabin, A. (2021). Features of Physical Rehabilitation of Volleyball Players with Patellar Tendinitis. *Human Sport Medicine*, 21(S1), 117-124.
- [6] Briner, W. W., Jr, & Benjamin, H. J. (1999). Volleyball injuries: managing acute and overuse disorders. *The Physician and sportsmedicine*, 27(3), 48–60. <https://doi.org/10.3810/psm.1999.03.720>
- [7] Ben Fredj, A., Rbai, H., Chatbouri, F., Berriri, M., Daadoucha, A., & Boughattas, A. (2023). Interest of longitudinal patellar tunnels in reconstructing chronic patellar tendon rupture with

- semitendinosus autograft: A case report. *Trauma case reports*, 49, 100969. <https://doi.org/10.1016/j.tcr.2023.100969>
- [8] Morelli, V., & Rowe, R. H. (2004). Patellar tendonitis and patellar dislocations. *Primary care*, 31(4), 909–ix. <https://doi.org/10.1016/j.pop.2004.07.003>
- [9] Fanelli, G. C., Stannard, J. P., Stuart, M. J., MacDonald, P. B., Marx, R. G., Whelan, D. B., Boyd, J. L., & Levy, B. A. (2010). Management of complex knee ligament injuries. *The Journal of bone and joint surgery. American volume*, 92(12), 2235–2246.
- [10] Andrew, S. (2011). Diagnostic Imaging Musculoskeletal: Trauma. *Radiology*, 2, 392.
- [11] Cucurulo, T., Louis, M. L., Thaumat, M., & Franceschi, J. P. (2009). Surgical treatment of patellar tendinopathy in athletes. A retrospective multicentric study. *Orthopaedics & traumatology, surgery & research : OTSR*, 95(8 Suppl 1), S78–S84. <https://doi.org/10.1016/j.otsr.2009.09.005>
- [12] Julie, K., Thomas, L., Eric, W., & Aly, T. (2000). Comparison of Rehabilitation Methods in the Treatment of Patellar Tendinitis. *Journal of Sport Rehabilitation*, 4, 304-314.
- [13] Griffiths, G. P., & Selesnick, F. H. (1998). Operative treatment and arthroscopic findings in chronic patellar tendinitis. *Arthroscopy*, 14(8), 836–839.
- [14] Aminaka, N., Pietrosimone, B. G., Armstrong, C. W., Meszaros, A., & Gribble, P. A. (2011). Patellofemoral pain syndrome alters neuromuscular control and kinetics during stair ambulation. *Journal of electromyography and kinesiology : official journal of the International Society of Electrophysiological Kinesiology*, 21(4), 645–651. <https://doi.org/10.1016/j.jelekin.2011.03.007>
- [15] Li, H., Cao, T., Zhou, H., & Hou, Y. (2019). Lip position analysis of young women with different skeletal patterns during posed smiling using 3-dimensional stereophotogrammetry. *American journal of orthodontics and dentofacial orthopedics*, 155(1), 64–70. <https://doi.org/10.1016/j.ajodo.2018.03.014>