

Comprehensive Diagnostic Approaches for Complex Tibiofibular Fractures

Shangkai Li

Qingdao No.58 High School of Shandong Province, Qingdao, China
gaimei@asu.edu.pl

Abstract. Tibiofibular fractures are often complex and involve both bone and soft tissue structures, making accurate diagnosis essential for appropriate surgical planning and better recovery outcomes. This paper explores current diagnostic techniques of complex tibiofibular fractures, particularly focusing on the application and advancement of CT and MRI imaging. The results of this paper show that conventional X-rays have significant shortcomings in terms of diagnostic sensitivity and accuracy. Multidetector computed tomography (MDCT) with three-dimensional (3D) reconstruction significantly improves fracture classification and spatial visualization, whereas magnetic resonance imaging (MRI) enhances soft-tissue and ligament assessment, further illustrating the importance of multimodal diagnostic planning for complex tibiofibular fractures. In addition, artificial intelligence-based image interpretation and CT-MRI fusion techniques show potential for improving diagnostic accuracy, speeding up processing, and evaluating outcomes. By evaluating the advantages and limitations of current techniques, this paper aims to provide a foundation for future developments in the precise and efficient diagnosis of complex tibiofibular fractures.

Keywords: Complex tibiofibular fractures, traditional X-ray, multidetector CT, 3D reconstruction, MRI.

1. Introduction

Complex tibiofibular fractures are among the most troubling orthopedic injuries to diagnose and manage in sports medicine, especially when involving open fracture, comminuted fracture, or intra-articular extension. Accurate and precise imaging is crucial throughout the whole process of treatment, from initial diagnosis and surgical planning to postoperative examination and rehabilitation. Although traditional X-ray imaging, which is widely available in hospitals, is effective and efficient, it often fails to display details like fracture lines or ligament structures. Thus, occult fractures and soft tissue damage may not be discovered by X-ray images. In response, computed tomography (CT) and magnetic resonance imaging (MRI) have emerged as essential diagnostic tools in diagnosing and managing these complex injuries.

CT has become the primary tool for analyzing bone structure, providing high-resolution images that improve doctors' understanding of fractures and aid in 3D reconstruction. CT imaging is very important for preoperative planning, as it can display the exact location of small bone fragments,

and doctors can also use CT to better align the fracture site, resulting in faster healing after treatment. MRI also plays a crucial role as it provides a visual representation of ligaments, soft tissues, and cartilages, which is particularly important in detecting occult fractures, evaluating postoperative healing, and observing the recovery of soft tissues and ligaments after surgery. The combination of CT and MRI greatly improves the accuracy of diagnosis. According to a research paper of CT-MRI combining diagnosis on hip injuries, CT and MRI techniques had 65.5% and 74.1% positive rates, respectively, while the positive rate of combining diagnosis of these two was 91.4%, which is huge progress compared to individuals [1]. Such a combination of diagnosing techniques is called multimodal diagnosis.

A perfect instance proving the advantages of multimodal diagnosis is the 2014 injury of NBA all-star player Paul George. Paul George suffered a complex tibiofibular fracture in a scrimmage for the USA team. His improper landing posture resulted in an open fracture of his right lower leg, accompanied by serious muscle and ligament tears. Doctors developed a surgical and rehabilitation plan for him based on CT and MRI scans. George has had a successful elite-level return-to-play supported by the powerful medical technology and his own efforts [2]. This illustrates how essential multimodal diagnosis is when facing complex tibiofibular fractures.

However, CT and MRI are often used alone due to resource limitations and time constraints. Emerging technologies, such as CT-MRI fusion and AI-auxiliary recognition, promise a prospect of diagnostic technology in the future. Inspired by Paul George's case, this paper analyzes various diagnostic tools, and it aims to support developing more effective and more accurate techniques that optimize the treatment process from injury diagnosis through to rehabilitation, and ultimately enhance performance for patients recovering from complex tibiofibular fractures, especially for athletes.

2. Clinical diagnostic challenges and needs of tibiofibular fractures

2.1. Classification of tibiofibular fractures

Tibiofibular fractures include a variety of injury patterns and clinical conditions. Effective diagnosis and management depend not only on examining the presence of a fracture, but also on classifying its type, evaluating its severity, and detecting soft tissue damage. A comprehensive classification system consists of multiple dimensions: whether to pierce the skin or not, soft tissue involvement, fracture morphology, and joint stability.

Tibiofibular fractures can be classified in terms of whether to pierce the skin or not, which is also known as an open/closed fracture. Open fractures are more severe, as the infection risk increases and complications are more likely to appear, while closed fractures are easier to heal. This perspective helps clinicians decide whether or not to take a surgery and inject antibiotics. Additionally, tibiofibular fractures are also described by their morphology, including transverse, angled, spiral, comminuted (broken into small fragments), or avulsion fractures. This type of classification can explain the injury mechanism of patients and determine the form of fixation imposed on the bone. Furthermore, tibiofibular fractures can be sorted based on their location, including fractures in the proximal 1/3, middle 1/3, and distal 1/3 of the tibia. In this classification method, different types of fractures can inform doctors of postoperative complications, such as osteofascial compartment syndrome. All fracture types in various classification methods may occur in tibiofibular fractures, which is why they are called complex tibiofibular fractures.

2.2. Requirements of imaging techniques and defects of traditional X-ray imaging

Clinicians hope to observe not only the details of the fractures and fracture lines for treatment planning, but also the presence of soft tissue and ligament damage or intra-articular extension, for assessing the condition of the injured. Misdiagnosis may cause improper treatment, persistent pain, or long-term effects.

Although X-ray imaging remains the main diagnostic technique due to its efficiency and availability, it has inevitable limitations. Standard 2D radiographs might fail to capture the details of tibiofibular fractures, such as slight displacement, soft tissue damage, or ligament involvement. A study on spiral tibial shaft fractures that may involve posterior malleolus (PM) injury shows the shortcomings of X-ray. In this study, 50 patients with tibial fractures took both X-ray and CT imaging, among whom 25 were determined to have additional PM fractures by CT. However, the detection rate of PM fracture was only 36.5% on average through a single 2D X-ray imaging technique. The more worrying thing is that in a group of clinicians who were not aware of the high incidence rate of PM fractures, only 17% of the injuries were identified by them. Even if another group of doctors were reminded to be careful with PM injuries, the detection rate improved to only 56%, and there were still nearly half of the injuries missed. In table 1, “PM fractures diagnosed” refers to the 25 patients who were determined to have a PM fracture in the CT imaging. “False positives” refers to the 25 patients who were not detected to have a PM fracture. The p-values are all lower than the significance level. The result clearly demonstrates that X-ray imaging lacks the sensitivity required for accurate diagnosis of spiral tibiofibular fractures [3].

Table 1. PM fractures diagnosed by traditional X-ray [3]

	Group 1(Not aware of high PM fracture incidence)	Group 2(Aware of high PM fracture incidence)
PM fractures diagnosed	4.25±3.77/25	14±0/25
False positives	0.5±0.5/25	2.5±1, 29/25
Sensitivity for X-ray	0.17	0.56
Specificity for X-ray	0.98	0.90




Figure 1. The comparison between X-ray images and CT images of the posterior malleolus fracture [3]

The comparison between X-ray images and CT images is shown from top to bottom in Figure 1. The image results indicate that, due to the fact that X-rays are 2D images, PM structure is obstructed by other bones and ligaments, making it difficult to distinguish whether there is a fracture. However, the presence of fracture lines can be seen in CT images. Another research study enrolled 67 patients who had an ankle joint problem. They did both X-ray and CT in three views and clinicians needed to assess whether there was a fracture without observing the CT images. Their findings, compared with CT findings, had an about 40% misdiagnosis rate. Thus, the researchers suggested that X-ray has a much lower accuracy for complex ankle injuries [4]. The ankle joint is located at the end of the tibia and is connected to the tibial condyle, so the study of ankle fractures applies to tibiofibular fractures.

2.3. Analysis and suggestion based on the case of Paul George

The injury to Paul George, which is a severe tibiofibular open fracture on his right lower leg, demonstrates the importance of a comprehensive and accurate diagnosis. Although specific images were not published, the complexity and severity of the injury, indicated by obvious bone deformation of his shank, likely require CT imaging for bone 3D reconstruction and MRI imaging for ligament and soft tissue evaluation. A single traditional X-ray image cannot provide enough information and conditions for doctors to make treatment plans and rehabilitation protocols. After such a devastating injury, Paul George could still get to a highly competitive level. This case proves that multimodal imaging weighs a lot in diagnosis, treatment, and readiness of return-to-play of athletes. In conclusion, tibiofibular fractures require precise classification, accurate diagnosis, and exact assessment of outcomes. The limitations of X-ray, especially in detecting complex or hidden injuries, emphasize the need for more advanced techniques like CT and MRI. The case of Paul

George reinforces this need, exemplifying how multimodal imaging supports successful recovery from serious orthopedic trauma.

3. Clinical value of CT technique

3.1. Accuracy of multidetector CT in the diagnosis of tibiofibular fracture

Nowadays, the CT technique has become a popular and advanced field of research. With the progress of technology, many different types of CT have been developed, which are classified by the locations that need to be scanned and by different scan methods (e.g., dual-energy CT, multidetector CT, cone-beam CT, etc.). For complex tibiofibular fractures, multidetector CT (MDCT) is the most commonly used. Unlike previous CT systems, MDCT can obtain precise and thin slices that are less than 1mm. This helps achieve rapid scan of volume, isotropic resolution, and multiplanar reformats. Clinicians could analyze the bone structure from the view of sagittal, transverse, and coronal planes.

A study done with 20 patients who had pelvic fractures shows that MDCT has a 100% accuracy in the diagnosis of pelvic fracture (See table 2). This research used a 16-layer scanner and 1mm slice thickness to take 3D and multi-planar reconstruction. What should be noticed is that in the figure (See Figure 2) below, the image on the left is a pelvic bone under X-ray, while the image on the right is a photo of a pelvic bone taken by MDCT. Posterior acetabular wall fracture was not detected by traditional X-ray at all. Instead, MDCT could clearly display that some small fragments were not being at their position [5].

Table 2. Validity of MDCT in detecting pelvic fracture [5]

Diagnostic Parameter	Value (%)
Sensitivity	100.0
Specificity	100.0
Positive Predictive Value (PPV)	100.0
Negative Predictive Value (NPV)	100.0
Accuracy	100.0



Figure 2. The comparison between X-ray images and MDCT images of the pelvic fracture [5]

3.2. Application of 3D reconstruction

3D reconstruction is the process that converts several 2D image slices into a 3D anatomical model. It plays a crucial role in diagnosis, surgical planning, and operations. The 3D model can be

displayed in various brightness, transparency, color, and frame, and it is able to be rotated, cut, and measured from any angle. A recent study used MDCT and CBCT of 24 dry skulls to test the accuracy and reliability of 3D reconstruction. The researchers used calipers measuring the size of the skulls 21 times directly, and these data would be compared with the ones generated by the 3D reconstruction model. As a result, there was no statistical significance between the value of direct measurements and 3D model measurements, showing an outstanding accuracy. Table 3 below provides data proving that MDCT makes accurate and precise anatomical measurements, which supports the application of 3D reconstruction models in clinical settings. Here, only abbreviations of locations are presented for a more direct and easier understanding. No statistically significant differences were found between direct and MDCT-based measurements for all anatomical landmarks (P-value > 0.05). Pelvic fractures are often complex, including tear fractures and open fractures, which are very similar to tibiofibular fractures. Therefore, even though this study mainly focused on the skull, the findings strongly agree in adopting MDCT 3D reconstruction to solve tibiofibular fractures, for a better treatment plan and assessment [6].

Table 3. The comparison of direct measurement and 3D model measurement by MDCT [6]

Landmark	Direct Measurement (mm)	MDCT 3D Reconstruction Measurement (mm)	p-value
Ba-Na	99.07±4.59	99.17±4.63	0.938
Ba-ANS	90.17±4.79	90.07±4.99	0.943
Na-A	56.37±2.98	56.50±2.93	0.879
Na-Me	120.19±5.50	120.21±5.37	0.991
Go-Co (Lt)	55.71±4.49	55.63±4.55	0.952

3.3. Analysis and suggestion on the application of CT technique in tibiofibular fractures

Computed Tomography (CT) has become the cornerstone of assessing and treating complex tibiofibular fractures, and has significant advantages. MDCT, which is one of the common types of CT, enables clinicians to visualize indistinct fracture lines, evaluate articular surface involvement, and determine the spatial orientation of bone fragments via its ability to print high-resolution and multi-planar images. Besides basic CT technique, 3D reconstruction is also a powerful tool for doctors to make surgical plans. Instead of a plain image from a certain angle, 3D models created by CT offer doctors a complete anatomical structure, with a study exemplified by this case study. In 2015, Dodd et al. used 3D reconstruction for a tibial plateau fracture's preoperative planning, which is also a kind of tibiofibular fracture. 3D reconstruction improved surgeons' spatial perception and confidence [7]. CT also benefits professional athletes. Paul George's successful surgery was to some extent achieved through precise and advanced CT technology. MDCT and 3D reconstruction are strongly recommended in complex tibiofibular fractures, especially open fractures and comminuted fractures. The use of CT in these settings improves both diagnostic accuracy and efficiency, contributing to better outcomes for patients.

4. Clinical value of MRI technique

4.1. Function of MRI

Magnetic resonance imaging (MRI) plays an important supplementary role in the process of treating a patient with a complex tibiofibular fracture. It is the best choice for assessing ligament, soft tissue,

and cartilage damage, which might be overlooked or underestimated under CT/X-ray. In addition, MRI is often used to exclude the possibility of other diseases and identify some missed and misdiagnosed cases. A study on finding the superiority of MRI for evaluation of sacral insufficiency fracture (SIF) used MRI, CT, and X-ray, respectively, to detect 77 patients. SIF is an occult fracture that is not easy to discover. Once again, it should be noted that the MRI scanning of the spine is essentially consistent with the scanning of the lower leg, so the interpretation of this study still has profound significance for tibiofibular fractures. What this research found is that MRI had a 100% sensitivity in this injury, while this value for CT and X-ray was 94.2% and 28.5%, worse than MRI. Table 4 illustrates the data in this study. In conclusion, MRI is a superior diagnostic tool for occult fracture [8].

Table 4. SIF diagnosis sensitivity of three techniques [8]

Technique	Sensitivity	p-value
X-ray	28.5%	P<0.01 vs. CT & MRI
CT	94.2%	P<0.01 vs. X-ray; p<0.05 vs. MRI
MRI	100%	P<0.05 vs. CT

4.2. Analysis and suggestion on the application of MRI technique in tibiofibular fractures

Although MRI is not a primary tool for diagnosis, it is considered as an effective supplement to the CT imaging for its various advantages mentioned above. Particularly, it is adopted to address joint instability or postoperative complications such as joint effusion. Gorelik et al. did research to support the long-term utility of MRI. In the systematic review of postoperative MRI surveillance of musculoskeletal injuries, MRI was proven to effectively detect early soft tissues' pathological changes, emphasizing its advantages in monitoring healing and identifying subclinical abnormalities [9]. This finding could be transferred to orthopedic traumas (complex tibiofibular fractures), where MRI helps with discovering ligament scarring and slow recovery of soft tissues, etc. The open fracture of Paul George must be accompanied by soft tissues as well, and MRI could be used selectively in such cases with suspected soft tissue damage and postoperative complications during rehabilitation.

5. Prospect of AI and multimodal diagnostic technology

With continued advancements in Radiography, the diagnosis of fractures, especially complex tibiofibular fractures and such difficult bone fractures, is developing towards greater precision and accuracy, higher efficiency, and better integration. Traditional X-ray, CT, and MRI all have their benefits, but emerging innovations advocate combining their advantages at the same time minimizing their disadvantages. Aayushi et al. reviewed 25 AI models that were used to classify and detect fractures. Among them, convolutional neural networks (CNN) performed with an accuracy ranging from 89% to 97%. Additionally, in some studies, AI even outperformed rookie doctors, and it matched the performance of experienced clinicians. The average dealing time was less than 10 seconds per case, which does provide a lot of help in urgent trauma conditions. A deep learning model has been trained for more than 10,000 images. Its sensitivity could reach 96.2%, and 93.4% for specificity. For complex tibiofibular fractures, an AI model can do AO/OTA classifications, reducing decision-making time [10]. Another area of progress is CT-MRI fusion technology, which is a type of multimodal technique. It combines bone images with high resolution (CT) and vision of

soft tissues (MRI) in a single workflow. In the clinical research of femoral neck injury patients, CT-MRI fusion technique enabled better localization of bone fragments and a complete visualization of surrounding tissues, far superior to using either of the two alone. The researchers noted that CT-MRI fusion imaging reduced diagnostic ambiguity and improved overall accuracy in clinical settings [1]. It is also valid in complex tibiofibular fractures, as bone fragments and ligament injuries often co-exist in such conditions. A combination of artificial intelligence and multimodal imaging has further potential. Future systems may integrate data from CT and MRI into AI models, and automated 3D construction and mock operations will presumably be invented to improve assessment of risk and outcomes. All these emerging technologies will have significant breakthroughs in efficiency, speed, accuracy, precision, and safety, which will have a profound impact on future medicine and might even cause a medical revolution.

6. Conclusion

This paper has evaluated the diagnostic tools that are currently used in managing tibiofibular fractures, focusing on the role of traditional X-ray, multidetector CT, 3D reconstruction, and MRI. According to the research, traditional radiography has defects to some extent, while it is money and time-saving. MDCT, especially when paired with 3D reconstruction, enhances the accuracy of fracture classification and perfects the treatment planning by providing a detailed vision of bone fragment displacement, fracture lines, and articular surface involvement. MRI, which helps with diagnosis, offers high sensitivity in evaluating ligament, cartilage, and soft tissue damage. These technologies complement each other and provide excellent advice for preoperative diagnosis, treatment planning, and postoperative rehabilitation assessment of patients, which will help athletes get a satisfying outcome and maintain a good condition when they return to the play. The results suggest that CT and MRI techniques can reduce missed diagnoses and misdiagnoses, enhance surgical precision, and shorten recovery time. In addition, the paper highlights the significance of multimodal imaging through the example (Paul George's case) of professional sports injury management, where accuracy and rehabilitation timelines are critical due to the complexity and severity of the injuries. Besides, emerging technologies such as AI-auxiliary interpretation of images and CT-MRI fusion offer further hope. AI detection has also demonstrated accuracy comparable to experienced doctors in identifying fracture types and processing images in a very short time. Moreover, CT-MRI fusion has shown potential in combining bone and soft tissue insights into a single diagnostic workflow, improving comprehensiveness and efficiency. Looking into the future, diagnostic imaging in orthopedic injuries is moving toward greater intelligence, integration, and personalization. These technologies are expected to play a role for clinicians not only in identifying injuries, but also in simulating operative approaches, predicting complications, and making rehabilitation plans. The combination of AI and multimodal imaging shows the future direction of orthopedic diagnostics. Continued research, interdisciplinary collaboration, and persistent experiments will be crucial to put innovation into practice, which will definitely promote the development of medical imaging.

References

- [1] Chen, J., Zheng, S., Song, Q., & Liu, A. (2020). Diagnostic Value of Magnetic Resonance (MR) Combined with Computed Tomography (CT) in Patients with Stress Injury of Femoral Neck. *Medical Science Monitor : International Medical Journal of Experimental and Clinical Research*, 26, e922170-1. <https://doi.org/10.12659/MSM.922170>

- [2] Sahisten, M., & Raj, S. (2025). Evaluating the impact of leg fractures on NBA players performance: a comparative study of pre-and post-Injury metrics / (pp. 9-10).
- [3] Lisitano, L., Röttinger, T., Wiedl, A., Rau, K., Helling, S., Cifuentes, J., Jehs, B., Härting, M., Feitelson, L. M., Gleich, J., Kiesl, S., Pfeufer, D., Neuerburg, C., Mayr, E., & Förch, S. (2023). Plain X-ray is insufficient for correct diagnosis of tibial shaft spiral fractures: a prospective trial. *European Journal of Trauma and Emergency Surgery*, 49(6), 2339–2345. <https://doi.org/10.1007/S00068-023-02285-X/FIGURES/1>
- [4] Szymański, T., & Zdanowicz, U. (2022). Comparison of routine computed tomography and plain X-ray imaging for malleolar fractures—How much do we miss? *Foot and Ankle Surgery*, 28(2), 263-268. <https://doi.org/10.1016/J.FAS.2021.03.025>
- [5] Housseini, A., Yousef, K. A., Al-Kady, L. M., & Elsherif, M. A. (2022). Role of Multidetector Computed Tomography (MDCT) in the Evaluation of Patients with Pelvic Trauma. *Suez Canal University Medical Journal*, 25(4), 12-18. <https://doi.org/10.21608/SCUMJ.2022.270482>
- [6] Kim, M., Huh, K.-H., YI, W.-J., Heo, M.-S., Lee, S.-S., & Choi, S.-C. (2012). Evaluation of accuracy of 3D reconstruction images using multi-detector CT and cone-beam CT. *Imaging Science in Dentistry*, 42(1), 25. <https://doi.org/10.5624/ISD.2012.42.1.25>, ”)
- [7] Dodd, A., Oddone Paolucci, E., & Korley, R. (2015). The effect of three-dimensional computed tomography reconstructions on preoperative planning of tibial plateau fractures: A case-control series. *BMC Musculoskeletal Disorders*, 16(1), 1-6. <https://doi.org/10.1186/S12891-015-0608-0/FIGURES/2>
- [8] Yamauchi, T., Sharma, S., Chandra, S., Tanaka, M., Fujiwara, Y., Arataki, S., Sharma, A., Yokoyama, Y., Oomori, T., Kanamaru, A., Masuda, S., Shimizu, N., Torigoe, K., & Honda, O. (2022). Superiority of MRI for Evaluation of Sacral Insufficiency Fracture. *Journal of Clinical Medicine* 2022, Vol. 11, Page 4968, 11(17), 4968. <https://doi.org/10.3390/JCM11174968>
- [9] Gorelik, N., Paruthikunnan, S., Uppal, A., Ervin, A. M., Ramanakumar, A. V., Quaiattini, A., Brophy, J. M., & Gyftopoulos, S. (2023). Usefulness of MRI-Based Local Surveillance After Surgical Treatment of Musculoskeletal Soft-Tissue Sarcomas: A Systematic Review and Meta-Analysis. *American Journal of Roentgenology*, 220(6), 805-816. https://doi.org/10.2214/AJR.22.28865/SUPPL_FILE/22_28865_SUPPL.PDF
- [10] Bhatnagar, A., Kekatpure, A. L., Velagala, V. R., & Kekatpure, A. (2024). A Review on the Use of Artificial Intelligence in Fracture Detection. <https://doi.org/10.7759/cureus.58364>