

Study of the Dirichlet Integral and Its Convergence Properties

Junyi Shi

*Kokomo High School, Indiana, USA
jshi13@ivytech.edu*

Abstract. The Dirichlet integral shows that an oscillating series can be convergent even though it is not absolutely convergent. It is this that is analyzed in this paper, and it talks about the reason for the integral $\int_0^\infty \sin x/x dx$ which it is approaching a certain finite value. The key properties of the integrand are defined first, with the most conspicuous being the profile about the origin and the oscillatory profile at infinity. The convergence of the integral is then analyzed using the oscillatory profile method, and classical profile convergence methods, including Dirichlet profile. The common formula of oscillatory integral is derived by taking into account the addition of an auxiliary parameter to the integral, which can be computed. In addition, a special case of the Dirichlet integral is briefly touched upon. Finally, the importance of oscillatory integrals is explained by looking at some of the most important cases in mathematics, physics, and engineering, which demonstrates its use in areas outside of theoretical mathematics.

Keywords: Dirichlet integral, improper integral, conditional convergence, Fourier analysis, sinc function

1. Introduction

In mathematical analysis, improper integrals play an important role in understanding functions that are not integrable in the traditional sense. Among them, the Dirichlet integral is one of the most famous examples. It is defined as

$$I = \int_0^\infty \frac{\sin x}{x} dx \quad (1)$$

and its value is known to be $\pi/2$ [1]. This result is surprising because the function $\frac{\sin x}{x}$ oscillates infinitely and is not absolutely integrable over $[0, \infty)$. The Dirichlet integral is important because it demonstrates a key concept in analysis: conditional convergence. Unlike absolutely convergent integrals, where the integral of the absolute value is finite, the Dirichlet integral converges due to the cancellation of positive and negative areas. This makes it a valuable example for understanding how oscillatory behavior affects convergence [2].

In addition to its theoretical significance, the Dirichlet integral has many applications in applied mathematics and physics. It plays a central role in Fourier analysis, especially in the study of the

sinc function, which is widely used in signal processing and communication systems. It also appears in problems involving wave motion and harmonic analysis. Previous studies in mathematical analysis, such as those found in classical textbooks by Apostol and Rudin, have provided rigorous proofs of the Dirichlet integral and explored its properties. These works emphasize the importance of parameter methods and advanced calculus techniques in evaluating such integrals [3].

This paper aims to present a clear and accessible analysis of the Dirichlet integral. Section 2 discusses its convergence and provides a detailed method for evaluating it. Section 3 introduces its applications in mathematics and physics. Finally, Section 4 summarizes the main conclusions.

2. Calculation of the Dirichlet integral

2.1. Convergence analysis

The Dirichlet integral Eq. (1) serves as a fundamental example in real analysis illustrating conditional convergence. At first glance, this integral appears problematic because the function $\frac{\sin x}{x}$ does not decay sufficiently fast to guarantee absolute convergence. In fact, the integral $\int_0^\infty \left| \frac{\sin x}{x} \right| dx$ diverges. Therefore, one has no single point of origin and the integral in any neighborhood of zero converges.

For small values of x , $\sin x$ is going to be bouncing back and forth between -1 and 1, getting ever so closer to 0. It is a slow decay, but works well in conjunction with oscillations. They if separated by enough time contribute positively and negatively and cancel out [4]. As x increases, $\sin x$ continues to oscillate between -1 and 1, while $1/x$ gradually decreases toward zero. Although this decay is slow, it becomes effective when combined with oscillations. Positive and negative contributions cancel each other over successive intervals, leading to convergence.

A more formal analysis is obtained by partitioning the interval into segments of length π . On each interval $[n\pi, (n+1)\pi]$, the sine function maintains a constant sign. Therefore, the following estimate holds:

$$\int_{n\pi}^{(n+1)\pi} \left| \frac{\sin x}{x} \right| dx \leq \int_{n\pi}^{(n+1)\pi} \frac{1}{x} dx \quad (2)$$

The contribution of each interval is decreasing because of. Cancellation is, on the contrary, performed by reversing signs in a similar way to an alternating series. Says that it demonstrates that people have conditional convergence of the integral [5].

The Dirichlet test of improper integrals is also a consequence of the Dirichlet integral. The reason is that, it tends to a test that alternating oscillation is limited and then tends to a limit when multiplied to another monotonic function with limit 0 not necessarily everywhere. Conversely, one has in this case, $\sin x \in [-1, 1]$. f is decreasing monotonically on it and limit $f(t)$ as t tends to infinity. Both these conditions guarantees that the integral, satisfies the condition of Dirichlet test. In this way, people innocently lose some absolute convergence but people ensure that they converge (not so unexpected). This hence offers a more stringent account to the intuitive account above in terms of cancellation and also serves as an additional support to the application of classical convergence tests in real analysis.

2.2. Evaluation via parameter method

To evaluate the Dirichlet integral, an auxiliary function is introduced:

$$F(a) = \int_0^{\infty} e^{-ax} \frac{\sin x}{x} dx, \quad a > 0 \quad (3)$$

The exponential factor ensures absolute convergence, allowing differentiation under the integral sign. Differentiation yields:

$$F'(a) = - \int_0^{\infty} e^{-ax} \sin x dx \quad (4)$$

The resulting integral is evaluated as $F'(a) = -\frac{1}{1+a^2}$. Integrating with respect to a , the following expression is obtained $F(a) = C - \arctan(a)$, where C is a constant [6].

To determine C , the limit as $a \rightarrow \infty$ is considered. In this limit, the exponential term forces the integral to approach zero, so $F(a) \rightarrow 0$. Since $\arctan(a) \rightarrow \frac{\pi}{2}$, it follows that $C = \frac{\pi}{2}$. Therefore,

$$F(a) = \frac{\pi}{2} - \arctan(a) \quad (5)$$

Taking the limit as $a \rightarrow 0^+$, the original Dirichlet integral is obtained

$$\int_0^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}. \quad (6)$$

This observation confirms the fact that an oscillatory functional can be able to take a finite value even though it is not absolutely convergent.

The parameter method is an analytical tool that deserves to be mentioned since it is not limited to this problem. The technique of an auxiliary parameter (differentiating under the integral sign) can be used to simplify some complex integrals. It is common in higher mathematics, in particular in mathematical physics. Here, the exponential factor is a regularizing factor in the Dirichlet integral, which becomes convergent at some intermediate steps. Once the tedious calculations of the limit are done the parameter is eliminated. This also exposes the deeper properties of the integral, especially the one between the inverse trigonometric functions, in addition to helping in the evaluation [7].

A more general form of the Dirichlet integral is given by

$$I = \int_0^{\infty} \frac{\sin(ax)}{x} dx \quad (7)$$

The value of this integral depends on the parameter a , and the result is $\pi/2$ for $a > 0$, 0 for $a = 0$, and $-\pi/2$ for $a < 0$. This formula reflects the dependence of the integral on the sign of a .

and highlights the symmetry of the sine function.

3. Applications of the Dirichlet integral

3.1. Fourier analysis

Another significant area of Dirichlet integral is the Fourier analysis, especially how functions can be expressed as frequency content. The function $\text{sinc}(x) = \frac{\sin x}{x}$ is a natural property in the Fourier transform of rectangular functions, which are typically used to represent signals of finite length. This Dirichlet integral process is useful in the explanation of the smooth and wavy results of these transforms, and not sharp edges [8].

Moreover, the Dirichlet Integral is associated with the inversion formula of Fourier Transform. Sine functions integrals are met when a function is being represented on top of its frequency representation. The convergence of these integrals is required to be mathematically correct. Without such a convergence, many fundamental discoveries of harmonic analysis would be invalid.

Besides this, Dirichlet integral is also related to the Dirichlet kernel, used in the partial sums of Fourier series. This is an oscillatory property of the kernel like the Dirichlet integral, e.g. cancellation. The relations above show that oscillatory integrals are significant in the analysis of convergence problems in Fourier analysis [9].

3.2. Signal processing

The Dirichlet integral is closely related to the sampling theory of signal processing upon which communication and data transmission is based. The sampling theory states that a continuous signal can be recovered without any information loss as long as the signal is sampled at a high rate. In reconstruction, the sinc function plays the role of an interpolation kernel [10].

When a signal is uniformly sampled the original continuous signal can be recovered by using shifted versions of the sinc function. This is directly because of Dirichlet integral that indicates that the sinc function is the most ideal reconstructive function of a signal. Even in the non-holy cases of these properties, people can reconstruct signals beautifully.

Practical applications of the theory include audio processing, image compression and digital communication systems. One such area is in the digitization of analog sound where sampling and reconstruction is important. These have mathematical basis, based on Dirichlet integral and are thus one of the most important concepts in modern technology.

3.3. Physics and mathematical significance

The Dirichlet integral also finds another application in physics, in particular in the study of wave phenomena. The oscillatory functions are applied to the wave propagation in the space in the case. The other types of integrals of this type, i.e., of the Dirichlet type of integral, are applied to investigate constructive and destructive patterns of interference in waves [11].

Sine integrals are considered in optics to analyse the diffraction patterns. Having a screen behind a small slit experiment and a monochromatic light behind the experiment can produce a pattern which could be described by functions containing the ratio of sin to x. The central maximum of the pattern and the smaller side peaks can be explained by the Dirichlet integral. Often quantum mechanics uses the Dirichlet integral in oscillatory integrals, and its generalization, the Feynman-Kleissmann integral, in particular. Such integrals determine the probability amplitudes and the

behavior of the particles at a microscopic scale. Such integrals possess convergence properties which are significant to the fact that predictions of physics are well-posed.

In mathematical analysis, the Dirichlet integral has a greater role to play, irrespective of the context. It is a bald example of a conditional convergence that illustrates how convergence may occur by oscillatory action as opposed to the swift decay. Behind this is a hint at numerous delicate questions of analysis. The Dirichlet test is one of the traditional convergence tests which are shown in the integral. The tests give definite cutoffs to the determination of whether an integral converges, where standard methods are inapplicable. Thus, Dirichlet integral is frequently used to teach higher-level calculus.

The Dirichlet integration is not only important in its immediate applications. They may co-operate, and produce important and stable results, and this is one of the general principles of mathematics. The principle is used in most areas, such as in asymptotic analysis, distribution theory, and partial differential equations. In the recent literature oscillatory integrals have been used to model more complex phenomena and to find approximate solutions to challenging problems. This is exemplified by the Dirichlet integral which is very simple and elegant in its thought and is therefore a good point of departure to further research. Its impact is evidenced in the progress in theory and in practice, showing the continuing significance of the integral in mathematics as well as in science.

4. Conclusion

To summarize, Dirichlet integral is a good example of how oscillatory character may cause convergence even though a Dirichlet is an absolute integrable. The integral analysis of $\int_0^\infty \frac{\sin x}{x} dx$ can be used to prove the correspondence between slow decay and alternating signs can cancel to give a finite result. This is also a source of one of the key concepts of mathematical analysis convergence does not need to be fast, but can also be a structural property of the integrand. The rigorous and elegant way of getting the exact value of the Dirichlet integral is to introduce a parameter of auxiliary and differentiate under the integral. Not only does it generate a hint of a strong correspondence between analysis and trigonometric functions, but also the non-geometrical derivation of inadvertently. In addition to this, the Dirichlet integral also finds application in applied mathematics as well as in pure mathematics (Fourier analysis, signal processing and physics) and is also applied in applied mathematics. Its relation to the sinc and sampling theory proves that it is significant practically in the modern technology and its manifestation in the wave phenomena proves its importance physically. The analysis of Dirichlet integrals demonstrates the strength of analytic tools, and the usefulness of oscillatory functions. It is between theory and practice and provides helpful hints about both mathematical structure and physical reality.

References

- [1] Stenger, F. (2000). Summary of Sinc numerical methods. *Journal of Computational and Applied Mathematics*, 121(1-2), 379-420.
- [2] Kreyszig, E. (2022). *Advanced engineering mathematics* (11th ed.). Wiley.
- [3] Zygmund, A. (2002). *Trigonometric series* (Vol. 1). Cambridge University Press.
- [4] Sugihara, M., & Matsuo, T. (2004). Recent developments of the Sinc numerical methods. *Journal of Computational and Applied Mathematics*, 164, 673-689.
- [5] Dhaigude, R. M., Chesneau, C., & Bagul, Y. J. (2020). About trigonometric-polynomial bounds of Sinc function. *Mathematical Sciences and Applications E-Notes*, 8(1), 100-104.
- [6] Huang, W., Qin, X. J., Zhang, Q. Y., & Li, S. S. (2024). Several calculation methods of Dirichlet integral based on mathematical physics methods. *Journal of Heihe University*, 15(11), 183-184.

- [7] Xu, G. J. (2024). A study on a class of Dirichlet generalized integrals. *Journal of Henan Finance University (Natural Science Edition)*, 33(2), 12-16.
- [8] Wang, S. L., & Wei, J. T. (2022). A new calculation method of Dirichlet integral. *Studies in College Mathematics*, 25(3), 62-63.
- [9] Abel, U., & Kushnirevych, V. (2023). Sinc integrals revisited. *Mathematische Semesterberichte*. Advance online publication.
- [10] Qian, C., Chen, X.-D., & Malešević, B. (2021). Tighter bounds for the inequalities of Sinc function based on reparameterization. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas*, 116(1), Article 29.
- [11] Wang, F. S. (2018). A generalization of Dirichlet integral. *Journal of Neijiang Normal University*, 33(6), 59-61.