

From Random Walk to Einstein's Diffusion Equation: A Review of Gas Diffusion Theories

Jiarui Wang

*School of Physics, University of Massachusetts Amherst, Amherst, USA
jiarui.wang@umass.edu*

Abstract. The study of gas diffusion is foundational to statistical mechanics and thermodynamics. This paper traces the evolution of diffusion models from the empirical roots of Fick's laws, through the discrete random walk theory and Stokes' drag formulation, culminating in Einstein's diffusion equation. By connecting macroscopic diffusion coefficients with the microscopic, stochastic motion of particles influenced by dissipative frictional forces, Einstein demonstrated that diffusion is a rigorous statistical outcome rather than a mere phenomenological law. Furthermore, this review explores contemporary extensions of this classical linear framework into non-linear paradigms. Specifically, it analyzes how the generalized framework introduced by Boon and Lutsko extends Einstein's stochastic master equation to capture complex, density-dependent diffusivity and anomalous transport. Ultimately, this comprehensive review bridges microscopic dynamics and macroscopic transport, providing a unified analysis of gas diffusion across heterogeneous and non-equilibrium systems.

Keywords: Gas Diffusion, Random Walk, Einstein-Smoluchowski Relation, Nonlinear Diffusion, Statistical Mechanics

1. Introduction

Diffusion is the transport of matter via thermal motion, and it governs how macroscopic thermodynamic systems spontaneously evolve toward chemical and thermal equilibrium. In gas kinetics, this process dictates mass transport driven by the chaotic thermal velocity of individual molecules. Despite its ubiquity, unifying observable macroscopic diffusion with its microscopic origins required nearly a century of theoretical evolution. During the nineteenth century, early transport investigations remained fundamentally empirical. In 1855, Adolf Fick formulated his laws of diffusion, establishing a linear relationship between particle flux and concentration gradients [1]. While Fick's laws successfully codified macroscopic behavior, they were purely phenomenological; the diffusion coefficient was an empirical constant uncomputable from first principles, leaving a profound theoretical void regarding molecular trajectories. The missing link emerged at the turn of the twentieth century through stochastic modeling and statistical mechanics. The concept of a "random walk" provided a discrete-time stochastic framework to track particle displacement from continuous collisions, finding an ideal empirical testing ground in Brownian motion. In 1905 and 1924, Albert Einstein brilliantly resolved the dichotomy between macroscopic and microscopic

regimes [2, 3]. By balancing classical thermodynamics (osmotic pressure gradients) with fluid mechanics (Stokes' viscous drag), Einstein derived a formulation computing the diffusion coefficient directly from molecular parameters. This bridged the gap between macro-transport and micro-fluctuations, validating the reality of atoms and Ludwig Boltzmann's statistical mechanics. In modern physics, classical linear diffusion represents an idealized limit. Real-world environments—such as dense gas mixtures, granular media, and biological cells—exhibit transport where the diffusion coefficient is not constant but depends intricately on local particle density, structural constraints, and environmental variations. This necessitates a transition from linear models to sophisticated nonlinear diffusion equations [4].

2. Fick's law

The formal mathematical theory of diffusion began with the work of the German physiologist Adolf Fick in 1855 [1]. Fick was heavily inspired by the prior physical formulations of Thomas Graham, who conducted extensive quantitative experiments on the diffusion of gases and salts in solution, as well as the mathematical analogies established by Joseph Fourier in his analytical theory of heat conduction. Fick recognized that the transport of matter down a concentration gradient was physically analogous to the transport of thermal energy down a temperature gradient.

Fick's first law quantitatively relates the diffusive flux vector, $\vec{J}$, to the local spatial gradient of the particle concentration, c . In a simplified one-dimensional representation, the law is written as:

$$J = -D \frac{\partial c}{\partial x} \quad (1)$$

where J represents the net diffusive flux (the amount of substance or number of particles passing through a unit cross-sectional area per unit time), D is the diffusion coefficient (typically expressed in dimensions of m^2/s), and $\partial c / \partial x$ is the spatial gradient of concentration along the x -coordinate. The negative sign is a mathematical expression of the second law of thermodynamics, ensuring that mass transport occurs spontaneously from regions of high chemical potential (high concentration) to regions of low chemical potential (low concentration).

To account for time-dependent phenomena where concentrations evolve dynamically, Fick combined his first law with the universal principle of mass conservation. In the absence of nuclear or chemical reactions that create or destroy particles, the time rate of change of concentration within an infinitesimal volume element must equal the negative divergence of the flux through its boundaries ($\partial c / \partial t = -\partial J / \partial x$). Substituting Fick's first law into this continuity equation yields Fick's second law of diffusion, commonly referred to as the linear diffusion equation:

$$\frac{\partial c}{\partial t} = \frac{\partial}{\partial x} \left(D \frac{\partial c}{\partial x} \right) \quad (2)$$

Assuming the medium is isotropic and homogeneous, the diffusion coefficient D remains uniform throughout the spatial domain, allowing it to be factored out of the spatial derivative:

$$\frac{\partial c}{\partial t} = D \frac{\partial^2 c}{\partial x^2} \quad (3)$$

This second-order partial differential equation laid the foundation for mass transport theory, chemical engineering, and environmental physics for the subsequent century. Given specific initial conditions—such as an localized instantaneous point-source of gas represented by a Dirac delta

function, $c(x, 0) = N_0\delta(x)$ —the analytical solution to Fick's second law yields a shifting Gaussian distribution:

$$c(x, t) = \frac{N_0}{\sqrt{4\pi Dt}} \exp\left(-\frac{x^2}{4Dt}\right) \quad (4)$$

While Fick's laws provided an elegant and incredibly precise tool for engineering applications and macroscopic experimental physics, they suffered from a profound conceptual limitation. As an empirical result, Fick's framework could describe how a substance spread out over time, but it was fundamentally silent on why diffusion occurred at the molecular level. Within this classical paradigm, the diffusion coefficient D was treated as a phenomenological black box. It was a property that could only be observed and cataloged through macroscopic measurement during this stage, but it could not be derived, explained, or calculated from any underlying microscopic properties of the gas molecules, such as their molecular mass, diameter, mean free path, or thermal velocity.

Consequently, the essential problem of this equation centered entirely on the nature of D . Without a microscopic theory, it was impossible to predict how D would change if the molecular structure of the gas was modified, or if the gas was subjected to extreme temperatures and pressures. Thus, the subsequent models of diffusion shifted focus away from purely macroscopic, continuum descriptions, concentrating instead on microscopic and statistical perspectives of constituent molecules and particles.

3. Random walk and brownian motion

To clarify the microscopic mechanisms driving diffusion, statistical mechanics models individual particle trajectories as a sequence of discrete random events, formalized as the "random walk" [5, 6].

In a basic one-dimensional model, a particle starts at $x = 0$ and takes independent steps of fixed length Δx at regular intervals Δt . At each step, it has an equal probability ($p = q = 1/2$) of moving right ($+\Delta x$) or left ($-\Delta x$). Let $X_n = \sum_{i=1}^n \xi_i$ represent the particle's position after n steps, where ξ_i are independent random variables. Because the expectation $\langle \xi_i \rangle = 0$, the average displacement remains zero:

$$\langle X_n \rangle = 0 \quad (5)$$

However, the variance or mean square displacement (MSD) quantifies the physical spreading. Since the steps are statistically independent, the variance of the sum equals the sum of individual variances ($\langle \xi_i^2 \rangle = (\Delta x)^2$):

$$\langle X_n^2 \rangle = n(\Delta x)^2 \quad (6)$$

To transform this discrete formulation into a continuous physical model, elapsed time is introduced via $n = t/\Delta t$. Substituting this into the MSD equation yields:

$$\langle x^2(t) \rangle = \left(\frac{(\Delta x)^2}{\Delta t}\right)t \quad (7)$$

By defining the macroscopic diffusion coefficient as $D = \frac{(\Delta x)^2}{2\Delta t}$, the continuous-time expression becomes:

$$\langle x^2(t) \rangle = 2Dt \quad (8)$$

In a three-dimensional isotropic space, the independent spatial components sum to:

$$\langle r^2(t) \rangle = 6Dt \quad (9)$$

This linear scaling with time ($\langle r^2 \rangle \propto t$) is the definitive hallmark of classical diffusion, contrasting with ballistic motion where displacement scales with t^2 . The random walk model successfully demonstrated how macroscopic diffusion equations spontaneously emerge from underlying microscopic stochasticity.

4. Stokes' law and the einstein relation

Albert Einstein physically derived the diffusion coefficient by unifying macroscopic hydrodynamics with the molecular-kinetic theory of heat [2, 3]. He modeled a suspension of microscopic particles as a dynamic balance between two competing physical forces.

The first is a thermodynamic driving force. A spatial concentration gradient ($\partial c / \partial x$) generates an osmotic pressure ($\Pi = ck_B T$) gradient according to van 't Hoff's law. This pressure drop manifests as an osmotic force per unit volume driving the particles downward:

$$F_{\text{osmotic}} = -k_B T \frac{\partial c}{\partial x} \quad (10)$$

This motion is counteracted by a hydrodynamic resistance. For a spherical particle of radius r moving at velocity v through an incompressible fluid with dynamic viscosity η , the dissipative drag is governed by Stokes' Law [7]:

$$F_{\text{drag}} = 6\pi\eta r v \quad (11)$$

In a steady-state transport regime, these forces reach dynamic equilibrium. Equating the total osmotic force per unit volume to the collective viscous drag ($c \cdot F_{\text{drag}}$) yields:

$$-k_B T \frac{\partial c}{\partial x} = c(6\pi\eta r v) \quad (12)$$

Solving for the drift velocity v and substituting it into the macroscopic flux definition ($J = cv$), we obtain:

$$J = - \left(\frac{k_B T}{6\pi\eta r} \right) \frac{\partial c}{\partial x} \quad (13)$$

Comparing this thermodynamic outcome directly to Fick's First Law ($J = -D \frac{\partial c}{\partial x}$) reveals the Stokes-Einstein relation:

$$D = \frac{k_B T}{6\pi\eta r} \quad (14)$$

This discovery was groundbreaking. It established a direct link between macroscopic transport (D, η) and microscopic parameters (r, k_B). Furthermore, it allowed Jean Perrin to experimentally measure Avogadro's number, definitively proving the atomic nature of matter [8]. Finally, it served as the earliest foundational manifestation of the Fluctuation-Dissipation Theorem in statistical mechanics.

5. From linear to nonlinear diffusion

The defining characteristic of classical linear diffusion is that the diffusion coefficient D is treated as a spatial and temporal constant, entirely independent of particle concentration ($D \neq D(c)$). This framework assumes that individual particles undergo independent random walks without interacting with or altering the transport properties of one another.

However, real-world physical, biological, and engineering environments frequently depart from this idealized behavior. Non-linear diffusion occurs when the transport rate is heavily influenced by local conditions, transforming the governing relation into a density-dependent partial differential equation:

$$\frac{\partial c}{\partial t} = \frac{\partial}{\partial x} \left(D(c) \frac{\partial c}{\partial x} \right) \quad (15)$$

When $D(c)$ follows a power-law relationship ($D(c) = D_0 c^m$), this formulation is known as the Porous Medium Equation. It successfully models complex macroscopic phenomena such as the structural rearrangements in granular sandpile diffusion [9], volume exclusion and crowding effects in cellular migration [10], and the propagation of density waves in non-linear traffic flow modeling [11].

To establish a rigorous microscopic foundation for these observations, Boon and Lutsko proposed a seminal generalization of Einstein's original framework [4]. Instead of modifying Fick's laws macroscopically, they returned to the fundamental probabilistic master equation that tracks particle positions over time. They introduced a density-dependent transition probability, positing that a particle's likelihood of hopping to an adjacent site is conditioned by the local concentration of the medium (e.g., restricted movement in crowded zones). By executing a gradient expansion on this modified master equation, they derived a generalized non-linear diffusion equation [4]. This proved that non-linear transport parameters naturally emerge from underlying microscopic correlations, allowing modern physics to model highly complex, interacting, and non-equilibrium systems with structural rigor.

6. Conclusion

This study investigated the historical, mathematical, and physical evolution of gas diffusion models, tracing the development from Fick's phenomenological laws to modern non-linear stochastic frameworks. The findings indicate that the trajectory of diffusion theory represents a continuous shift from macroscopic empirical descriptions to microscopic statistical-mechanical rigor. By critically analyzing the conceptual linkages between Fick's laws, Markovian random walks, the Fluctuation-Dissipation theorem, and modern density-dependent generalizations, this research contributes to the existing body of knowledge by consolidating disparate transport physics frameworks into a cohesive lineage. These findings extend previous theories by demonstrating that macroscopic diffusion is a robust statistical consequence of underlying microscopic stochasticity and dissipative viscous forces. Practically, this study has significant implications for optimizing

industrial gas separation, predicting atmospheric pollutant dispersion, and modeling nutrient transport across crowded cellular membranes. However, this study is limited by its focus on idealized spherical particles in isotropic media and the absence of direct computational simulations, which may affect the generalizability of the findings to elongated molecules or multi-phase environments. Future study could focus on employing high-performance molecular dynamics simulations to precisely map the density thresholds where linear transport transitions into the non-linear paradigm. In the future, the author intends to develop a customized stochastic algorithm that integrates variable thermal environments into the density-dependent master equation. Overall, this study provides new insights into the enduring conceptual power of statistical mechanics, paving the way for advanced transport modeling in complex, non-equilibrium systems across physics, biology, and engineering disciplines.

References

- [1] Fick, A. (1855). Ueber Diffusion. *Annalen der Physik*, 170(1), 59-86. <https://doi.org/10.1002/andp.18551700105>
- [2] Einstein, A. (1905). Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen. *Annalen der Physik*, 322(8), 549-560. <https://doi.org/10.1002/andp.19053220806>
- [3] Einstein, A. (1924). The Stokes-Einstein Law for diffusion in solution. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 105(733), 412-416. <https://doi.org/10.1098/rspa.1924.0028>
- [4] Boon, J. P., & Lutsko, J. F. (2007). Nonlinear diffusion from Einstein's master equation. *EPL (Europhysics Letters)*, 80(6), 60006. <https://doi.org/10.1209/0295-5075/80/60006>
- [5] Pearson, K. (1905). The problem of the random walk. *Nature*, 72(1865), 294. <https://doi.org/10.1038/072294b0>
- [6] von Smoluchowski, M. (1906). Zur kinetischen Theorie der Brownschen Molekularbewegung und der Suspensionen. *Annalen der Physik*, 326(14), 756-780. <https://doi.org/10.1002/andp.19063261405>
- [7] Stokes, G. G. (1851). On the effect of the internal friction of fluids on the motion of pendulums. *Transactions of the Cambridge Philosophical Society*, 9, 8-106.
- [8] Perrin, J. (1909). Mouvement brownien et réalité moléculaire. *Annales de Chimie et de Physique*, 18, 5-114.
- [9] de Gennes, P. G. (1999). Granular matter: a tissue of problems. *Acta Materialia*, 47(15), 4137-4146. [https://doi.org/10.1016/S1359-6454\(99\)00294-0](https://doi.org/10.1016/S1359-6454(99)00294-0)
- [10] Murray, J. D. (2002). *Mathematical Biology: I. An Introduction* (3rd ed.). Springer-Verlag.
- [11] Whitham, G. B. (1974). *Linear and Nonlinear Waves*. John Wiley & Sons. <https://doi.org/10.1002/9781118032954>